UMASS/AMHERST
312066 0311 0332 3

MASS
EA
31
.2:M
382/3/

University of
Massachusetts
Amherst

LIBRARY

Digitized by the Internet Archive
in 2013 with funding from
Boston Library Consortium Member Libraries

http://archive.org/details/massachusettscoa1999mass

MASS. EA31.2: M382/3/999

MASSACHUSETTS COASTAL COMMERCIAL LOBSTER TRAP SAMPLING PROGRAM MAY-NOVEMBER, 1999

Bruce T. Estrella
and
Robert P. Glenn

GOVERNMENT DOCUMENTS
COLLECTION

NOV 2 0 2000

University of Massachusetts
Depository Copy

COMMONWEALTH OF MASSACHUSETTS
Division of Marine Fisheries
Paul J. Diodati, Director

Department of Fisheries, Wildlife and
Environmental Law Enforcement
David M. Peters, Commissioner

Executive Office of Environmental Affairs
Robert Durand, Secretary
August 28, 2000

TABLE OF CONTENTS

ABSTRACT	ii
INTRODUCTION	1
STUDY AREA	1
SAMPLING PROCEDURE	1
ANALYTICAL PROCEDURES	3
RESULTS AND DISCUSSION	4
Commercial Lobster Sampling	4
Index of Pre-Recruit Abundance	7
Water Temperature Time Series	10
ACKNOWLEDGEMENTS	14
LITERATURE CITED	15
APPENDIX	16

Publication No. 18458 - 28 - 208 - 10/00 - .97 - C.R.
Approved by: Philmore Anderson III, State Purchasing Agent

ABSTRACT

This is the Massachusetts Division of Marine Fisheries nineteenth annual assessment of the status of the American lobster resource in Massachusetts coastal waters. During the period of May through November, 1999, seventy-nine (79) sampling trips were made aboard commercial lobster vessels. A total of 31,637 lobster was sampled from 15,102 trap hauls. The catch rate of marketable lobster, 0.902 lobster per trap, was 36% higher than the 1998 index, 0.662. The proportion of females ovigerous, 17.4%, was higher than in the previous year (16.4%). The coastwide fishing mortality estimate, 1.22, was lower than the 1998 index of 1.36. Exploitation rate, 0.67, remained unchanged, while mean carapace length of marketable lobster, 89.6mm, and mean size of egg-bearing females, 88.1 mm, increased fractionally. The cull rate, 20.8%, was similar to the 1998 estimate of 20.9%. Less than 1% of the lobster sampled from traps were dead.

An index of pre-recruit abundance was created using data from our sea sampling database. The catch rate of lobster in the pre-recruit size class provided the basis of the index. The index was calculated from a multiple regression of log transformed catch rates with the following factors: month, year, lobsterman. The back-transformed regression coefficient associated with the factor "year", adjusted for the effects of the other analyzed factors, provided the index of pre-recruit abundance. The relationship between the index and Massachusetts territorial waters catch in the following year was modelled using a power function. The resulting equation was used to calculate predicted landings which differed from the actual landings for the years 1981-1999 by 0.43 to 25.54% with a mean difference of 8.65%.

A time series of data from our bottom water temperature monitoring program is presented for seven locations in Buzzards Bay, Cape Cod Bay, and Massachusetts Bay.

INTRODUCTION

This is the Massachusetts Division of Marine Fisheries (DMF) nineteenth annual assessment of the status of the American lobster resource in Massachusetts coastal waters. Since the lobster resource supports the most economically important single-species fishery in Massachusetts coastal waters, a long-term coastwide lobster monitoring program yielding biological and catch per unit effort data was devised and initiated in Massachusetts in May, 1981. A sea sampling/survey design was chosen by which both catch per unit effort and biological data could be collected temporally and areally with sufficient precision for stock assessments. The objective was to assess variations in population parameters due to environmental factors, fishing pressure, and regulatory changes.

Data collected during the 1999 coastwide commercial lobster trap sampling program are summarized below. Parameter trends occurring during the 1981-1999 study period are presented.

STUDY AREA

The study area is primarily defined by the Massachusetts territorial sea, except where lobstering activities of cooperating commercial lobstermen exceeded territorial boundaries (Figure 1). Territorial waters total 5,322 sq km (2,055 sq n mi), of which an estimated 60% is considered major lobster habitat. Six sampling regions, Cape Ann, Beverly-Salem, Boston Harbor, Cape Cod Bay, outer Cape Cod, and Buzzards Bay, were chosen for coverage of the major lobstering regions of the state. For convenience, these regions are depicted in Figure 1 as generalized hatch-marked areas wherein lobster gear sampled may be discontinuously distributed.

SAMPLING PROCEDURE

Sampling of coastal waters was accomplished by monitoring catches during the normal lobstering operations of volunteer commercial lobstermen in each designated region. Multiple lobstering operations were observed to reduce bias from varying degrees of lobstering skill and to enhance areal coverage. Pot-sampling trips were day trips, conducted a minimum of once per month per region during the major lobstering season, May-November.

Utilizing portable cassette tape recorders, sea samplers recorded carapace length (to the nearest mm); sex; and condition, including the degree of shell hardness, culls and other shell damage, external gross pathology, mortality, and presence of extruded ova on females (ovigerous). Catch in number of lobster, number of trap hauls, set-over-days, trap and bait type were also recorded. Trap locations were recorded from LORAN and plotted on nautical charts. Depth information was then acquired from the charts as a coastwide standard to avoid variability from tidal fluctuations.



Figure 1. Map of Massachusetts with six lobster sampling regions. Gray dots indicate location of trap trawls sampled in 1999, and dotted line represents the territorial sea boundary.

ANALYTICAL PROCEDURES

Data were computer coded and keypunched with a microcomputer data entry program. The data base was subsequently transferred for analysis to the Massachusetts Executive Office of Environmental Affairs' (EOEA) Digital Equipment Corporation VAX-11/780 computer system. A computer auditing process was used to uncover keypunch and recording errors and statistical analyses were performed with SPSS (Nie 1983) statistical sub-programs.

Because parameter means exhibit significant regional and monthly variation, an areal and temporal data weighting scheme was incorporated into analytical software. As a result, each month's data contribute equally to regional parameter means which are weighted by area in square nautical miles to generate coastwide means.

Unless specified otherwise, the terms "legal" or "legal sized" lobster include all lobster in the carapace length category $\geq$ 82.6 mm. The marketable segment of this category, which excludes ovigerous females, is analyzed separately and referred to as "marketable lobster". The sublegal length category includes all lobster < 82.6 mm.

The catch rates of marketable lobster are expressed as CTH'_3. This is catch per trap haul standardized to 3 set-over-days (Estrella and McKiernan 1989).

Estimates of total instantaneous mortality (Z) and total annual mortality ($A=1-e^{-Z}$) were computed by two methods which produce extremes in the possible range of estimates. The method of Gulland (1969) requires computation of the regression line slope of natural log-transformed numbers at estimated age (15% molt groups, 14% for Buzzards Bay, were derived from tagging data). Beverton and Holt's (1956) process employs von Bertalanffy Growth Equation parameters (from Fair 1977) and mean and minimum length of exploitable sizes.

Estimates of fishing mortality (F) were calculated with cohort analysis (Pope 1972, Jones 1974). Rates of exploitation were calculated with the equation $u=FA/Z$, where F= fishing mortality, A= total annual mortality, and Z= total instantaneous mortality.

Lobster landings data were derived from lobstermen's catch reports which are compiled annually by the DMF Commercial Fisheries Statistics Project.

Since current management strategy stresses uniform coastwide regulations, all data are grouped for a coastwide analysis. However, the uniqueness of the Massachusetts coastline, its role as a temperature barrier which profoundly affects many marine species (Colton 1964), and the influence of offshore lobster stocks on the inshore resource mandate a regional data treatment as well.

RESULTS AND DISCUSSION

Commercial Lobster Sampling

During the period of May through November, 1999, seventy-nine (79) sampling trips were made aboard commercial lobster vessels in Massachusetts coastal waters. A total of 31,637 lobster was sampled from 15,102 trap hauls.

The 1999 coastwide mean catch per unit effort index (CTH'_3), 0.902 marketable lobster per trap, was 36% higher than the 1998 index, 0.662 (Appendix Table 1). Total Massachusetts commercial landings, 15,869,927 lbs, increased by 20% from 1998. Landings from territorial waters, 10,066,631 lbs, increased by 31%. Landings and catch rate trends are depicted in Figure 2. The coastwide mean catch rates of sublegal lobster decreased by 27% (CTHSOD) and 23% (CTHAUL) from 1998. These represent new time series lows (Appendix Tables 2 and 3).

Increased legal catches and decreased sublegal catch rates may have been partly due to improved trap efficiency from the increase in escape vent size, effective 1 February, 1999 and good conditions for molting across the legal size threshold.



Figure 2. Catch per unit effort of marketable American lobster from commercial trap sampling and Massachusetts lobster landings from territorial water, 1981 - 1999.



Figure 3. Relative abundance of ovigerous female American lobster in percent total females and catch per effort, Massachusetts coastal waters, 1981 - 1999.

Of all females sampled during 1999, 17.4% were ovigerous compared to 16.4% in 1998 (Appendix Table 4). However, trends in statewide CPUE of ovigerous females (Appendix Tables 5-6) either increased or remained unchanged and are depicted in Figure 3.

Approximately 91% of the legal catch in our inshore regions (Cape Ann south through Cape Cod Bay and Buzzards Bay) was comprised of new recruits (83 mm-94 mm CL), i.e., lobster which recruited to the legal size range during their most recent molt (Appendix Table 7). This index of the effect of fishing pressure on the size frequency was similar to 1998. The index fluctuated from 58% in 1998 to 46% in 1999 for the primarily offshore migrant lobster sampled east of Cape Cod. Estimates of total mortality (Z) for inshore Gulf of Maine regions (Z = 1.14-3.14, A = 68%-96%) and Buzzards Bay (Z = 2.42-3.81, A = 91%-98%) depict a heavily exploited resource while those for the outer Cape Cod region (Z = 0.58-0.59, A = 44%-45%) indicate that a lower level of fishing pressure was exerted on this lobster group (Appendix Tables 8A and 8B).

Estimates of instantaneous fishing mortality (F), the proportion of all deaths which are attributed to fishing, ranged from 0.52 off outer Cape Cod to 2.14 in Buzzards Bay (Appendix Table 9A). Estimates of F using DeltaT values in place of von Bertalanffy growth parameters were lower (Table 9B), ranging from 0.39 off outer Cape Cod to 1.07 in Buzzards Bay. Exploitation rates (u), i.e. the fraction of the population that is removed by fishing, was unchanged from 1998 data (Appendix Table 10).

The relationship between fishing mortality, rate of exploitation, and mean lobster size is depicted in Figure 4. Carapace length exhibited a downward trend as fishing mortality and exploitation rates increased through 1987. Thereafter, increases in mean carapace length of 0.7

mm occurred in 1988 (mean size = 88.2 mm) and 1989 (mean size = 88.9 mm, Appendix Table 11) which reflected the similar numerical change in the minimum legal size during those years. Carapace length then fluctuated downward until 1994, and increased during 1995 – 1997, reaching a time series high in 1999. Fishing mortality rates for all regions combined edged upward to a time-series high of 1.48 in 1993, declined in 1994-1996 along with exploitation rates, then fluctuated upward in 1997-1998, before declining to a 1982 level in 1999. The relative change in size frequency between 1998 and 1999 is depicted by the overlay in Figure 5.

Sublegal sized lobster averaged 77.4 mm carapace length during 1999 compared to 77.2 mm during 1998 (Appendix Table 12). The mean size of all ovigerous females increased between 1998 (86.4 mm) and 1999 (88.1 mm).



Figure 4. Relationship between exploitation rate, fishing mortality, and mean carapace length of marketable American lobster, Massachusetts coastal water, 1981 - 1999.

The percentage of culls (lobster with one or both claws missing or regenerating) among all lobster sampled was similar between 1998 (20.9%) and 1999 (20.8%) (Appendix Table 14). The cull rates for legal, marketable, and sublegal size groups remained comparable between years, decreasing by only a fraction of a percent (Appendix Tables 15-17).



Figure 5. Length frequencies of trap-caught American lobster, Massachusetts coastal waters, 1998 - 1999.

The coastwide incidence of lobster found dead in traps was 0.18%. This was similar to that of the previous year (Appendix Table 18) and is acceptably low.

Index of Pre-Recruit Abundance

We created an index of abundance for pre-recruit lobster with data from our sea sampling program. The catch rate (number caught per trap haul) of lobster in the pre-recruit size class provided the basis of the index. The pre-recruit size range changed with increases in minimum legal size (68-80 mm carapace length for the years 1981-1987; 69-81 mm for 1988; and 70-82 mm for 1989 and later). The index was calculated from a multiple regression of logarithmically-transformed catch rates with the following factors: month, year, lobsterman. Other factors including soak time, bait, and trap type were incorporated in the preliminary regressions but failed to improve the fit and so they were excluded from the model. The back-transformed regression coefficients associated with the factor "year", adjusted for the effects of the other analyzed factors, provide the index of pre-recruit abundance (IPA). This approach using regression/ANOVA modelling in the standardization of catch rates has been used successfully to create indices of juvenile abundance in the Western Australia rock lobster (*Panulirus cygnus*) fishery (Caputi and Brown, 1986).

The relationship between the IPA and territorial catch in the following year (Figure 6) was modelled using a power function resulting in the following equation:

$$\mathbf{Catch}_{t+1} = 9{,}552{,}371(\mathbf{IPA}_t)^{0.671}$$

r^2=0.255; p=0.005

where $\mathbf{Catch}_{t+1}$ = territorial catch in year t+1

$\mathbf{IPA}_t$ = index of pre-recruit abundance in year t



Figure 6. Relationship between the index of pre-recruit abundance and Massachusetts territorial catch one year later.

This equation was used to predict territorial landings. The model predicts a decrease in territorial landings for 2000 to 8,363,955 lbs.

The relationship between predicted territorial landings and actual territorial landings is shown in Figure 7. The predicted values differed from the actual values by 0.43% to 25.54% with a mean difference of 8.65%. The relatively tight fit between predicted and actual landings (r=0.54, p=0.02) indicates the model provides modest predictive power for territorial landings.



Figure 7. Relationship between predicted landings based on the pre-recruit abundance index and actual territorial landings.

The model's predictive capabilities were evaluated via graphical analysis. The absolute percent change in actual landings from one year to the next was overlaid with the absolute percent change in predicted landings. Percent change of actual landings (3.21% to 31.41 %, mean = 11.13%) varied from year to year and had a higher mean percent change than predicted landings(0.23% to 18.6%, mean = 5.54%) (Figure 8). This indicates that the catch rate of pre-recruit lobsters in the previous year accounts for approximately 50% of the variation in territorial landings over the time series. The remaining variance associated with territorial landings is likely attributed to changes in fishing effort and environmental conditions, especially temperature. As is, the model has limited forecasting value. However, it is likely that its predictive capability would be enhanced if accurate estimates of changes in fishing effort and water temperature were integrated into the model.



Figure 8. Absolute percent change in MA territorial landings and predicted landings.

Water Temperature Time Series

In 1985, a coastal bottom water temperature monitoring project was initiated. Temperature monitors (RYAN Tempmentor) have been deployed for various lengths of time at several sites in Cape Cod Bay, outside Boston Harbor, and Buzzards Bay (Figure 8). Some of these sites are located on ship wrecks.

The longest time series of bottom temperatures is from Cleveland Light in Buzzards Bay. The last monitor to be deployed was at Rocky Point, off Plymouth. The Rocky Point, Manomet Point, *Endicott*, and *Mars* sites represent the 0-30 ft., 30-60 ft., 61-90 ft., and 91-120 ft. depth strata, respectively, in Cape Cod Bay. The *Romance* (off Boston Harbor), and Buzzards Bay-South sites are located at 70-80 ft. and provide data from the north-south extremes in our series. The Cleveland Light monitor is located in 30 feet of water.

Monitors are retrieved and replaced annually by divers. Although the time series contained data from seven monitors at one point, we currently collect data from only six sites because the monitor at the *Endicott* site was lost and has not been replaced. Figures 9 and 10 present the bottom water temperature at sites in Buzzards Bay and Cape Cod Bay/Massachusetts Bay, respectively. Figure 11 provides a comparison between the annual mean bottom temperature at Cleveland Light, Manomet, *Mars*, and *Romance* and the annual mean surface temperature at Boston and Woods Hole provided by NOAA/NOS.



Figure 9. Map of Massachusetts with approximate locations of seven bottom temperature monitors indicated.



Figure 10. Mean monthly bottom water temperatures at two sites in Buzzards Bay.



Figure 11. Mean monthly bottom water temperatures at five sites in the Gulf of Maine.



Figure 12. Mean annual bottom water temperatures at four sites monitored by the Coastal Lobster Project and mean annual surface temperature at two sites monitored by NOAA/NOS, 1986 - 1999.

ACKNOWLEDGEMENTS

We are indebted to the many commercial lobstermen whose cooperative spirit and concern for the American lobster resource sustain our lobster monitoring program. Gratitude is also extended to Bill Hoffman, Brad Chase, Rob Johnston, Matt Camisa, and Brian Kelly for data collection, Ann Spires, for data entry, and James Fair who administered the project and reviewed the manuscript. We also thank Thomas Hoopes for his data entry software design and assistance in data quality control.

LITERATURE CITED

Beverton, R. J. H. and S. J. Holt. 1956. A review of methods for estimating mortality rates in exploited populations, with special reference to sources of bias in catch sampling. Reports et Proces-Verbaux des Reunions 140:67-83.

Caputi, N., and R. S. Brown. 1986. Relationship between indices of juvenile abundance and recruitment in the western rock lobster (*Panulirus cygnus*) fishery. Can. J. Fish. Aquat. Sci. 43:2131-2139.

Colton, J. B. 1964. History of oceanography in the offshore waters of the Gulf of Maine. U.S. Fish and Wildlife Service, Special Scientific Report, Fisheries No. 496. 18p.

Estrella, B. T. and D. J. McKiernan. 1989. Catch-per-unit-effort and biological parameters from the Massachusetts coastal lobster (*Homarus americanus*) resource: description and trends. NOAA Tech. Rep. NMFS 81. 21 p.

Fair, J. J., Jr. 1977. Lobster investigations in management area I: Southern Gulf of Maine. NOAA, NMFS State-Federal Relationships Division, Mass. Lobster Report No. 8., April 21, 1975 - April 20, 1977, 8 p. and Appendix, 14 p.

Gulland, J. A. 1969. Manual of methods for fish stock assessment. Pt. 1, fish population analysis. FAO Manuals in Fisheries Science 4, 154 p.

Jones, R. 1974. Assessing the long term effects of changes in fishing effort and mesh size from length composition data. ICES CM 1974/F:33.

Nie, N. H. 1983. SPSS: statistical package for the social sciences, McGraw-Hill, New York, 806 p.

Pope, J. G. 1972. An investigation of the accuracy of virtual population analysis using Cohort analysis. Research Bulletin Int. Comm. NW Atlant. Fish. 9:65-74.

Sokal, R.R. and F.J. Rohlf. 1969. Biometry. W.H. Freeman and Company, San Francisco. 776 p.

APPENDIX

Table 1. CTH'3, by state and region, for all marketable lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.767	0.785	0.803	0.696	0.825	0.816	0.737	0.820	0.751	0.826	0.795	0.716	0.671	0.966	0.893	0.792	0.776	0.662	0.902
Cape Ann	0.732	0.808	0.624	0.663	0.634	0.699	0.669	0.496	0.721	0.904	0.868	0.724	0.770	1.015	0.979	1.272	1.106	0.645	0.699
Beverly-Salem	0.934	0.898	0.881	0.835	0.663	0.496	0.611	0.661	0.639	0.827	0.586	0.390	0.509	0.898	0.840	0.788	0.419	0.767	1.005
Boston Harbor	---	---	---	1.108	1.254	1.096	1.058	1.057	1.123	1.224	1.160	0.734	0.750	0.725	0.626	0.534	0.638	0.462	0.787
Cape Cod Bay	0.710	0.776	0.680	0.479	0.716	0.822	0.533	0.752	0.539	0.630	0.693	0.567	0.494	1.052	0.906	0.662	0.760	0.633	1.021
Outer Cape Cod	0.808	0.824	0.765	0.598	0.856	0.811	0.937	0.861	0.923	1.219	1.148	1.339	1.021	1.105	1.117	1.027	0.837	0.785	0.893
Buzzards Bay	0.611	0.571	1.110	0.870	0.953	0.907	0.952	1.064	0.934	0.598	0.575	0.817	0.834	0.852	0.893	0.866	1.169	0.719	0.740

Table 2. CTHSOD, by state and region, for all sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.580	0.672	0.718	0.521	0.647	0.700	0.578	0.509	0.695	0.716	0.665	0.465	0.542	0.402	0.410	0.343	0.363	0.295	0.215
Cape Ann	0.067	0.109	0.586	0.450	0.395	0.474	0.417	0.388	0.670	0.589	0.728	0.726	0.447	0.658	0.800	0.573	0.831	0.356	0.171
Beverly-Salem	0.708	0.711	1.263	0.948	0.833	0.801	0.863	0.353	0.780	0.408	0.324	0.411	0.406	0.314	0.334	0.277	0.231	0.364	0.233
Boston Harbor	---	---	---	0.901	1.162	1.138	1.156	0.639	0.966	1.103	0.924	0.839	0.800	0.690	0.782	0.688	0.651	0.754	0.508
Cape Cod Bay	0.710	1.013	0.639	0.322	0.594	0.551	0.371	0.438	0.595	0.727	0.716	0.298	0.436	0.313	0.307	0.217	0.256	0.204	0.162
Outer Cape Cod	0.037	0.024	0.038	0.033	0.035	0.027	0.088	0.064	0.066	0.078	0.077	0.088	0.075	0.045	0.060	0.051	0.053	0.043	0.025
Buzzards Bay	0.787	0.620	0.638	0.785	0.848	1.312	0.871	1.153	1.188	1.236	1.072	0.784	1.156	0.649	0.492	0.540	0.510	0.254	0.272

Table 3. CTHAUL, by state and region, for all sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	1.473	1.401	1.624	1.389	1.705	1.899	1.873	1.736	2.297	2.216	1.996	1.460	1.720	1.389	1.457	1.332	1.417	1.259	0.967
Cape Ann	0.256	0.199	1.044	0.909	1.031	1.126	1.143	1.062	1.765	1.782	1.783	1.661	1.562	1.725	2.323	1.660	1.940	1.074	0.606
Beverly-Salem	1.855	1.713	2.526	2.504	2.567	2.435	3.482	1.862	3.477	1.867	1.563	1.502	1.540	1.717	1.920	1.654	1.988	2.038	1.577
Boston Harbor	---	---	---	2.773	3.038	3.314	3.334	1.959	3.104	3.382	2.451	2.069	2.284	2.189	2.390	2.511	2.258	2.864	1.660
Cape Cod Bay	1.544	1.680	1.345	0.825	1.337	1.512	1.031	1.442	1.742	1.921	2.086	1.065	1.334	1.033	1.102	0.873	0.926	0.701	0.680
Outer Cape Cod	0.233	0.145	0.210	0.189	0.160	0.161	0.324	0.353	0.306	0.453	0.452	0.490	0.474	0.288	0.359	0.372	0.354	0.301	0.172
Buzzards Bay	2.381	1.916	2.316	1.965	2.452	3.118	3.090	3.722	3.984	3.994	3.181	2.602	3.501	2.179	1.599	1.835	2.051	1.546	1.456

Table 4. Percent of females ovigerous, by state and region, for all American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	5.9	7.7	10.9	9.1	8.6	9.1	9.2	8.8	10.0	10.9	9.8	11.9	14.3	10.7	14.1	15.4	14.7	16.4	17.4
Cape Ann	1.7	3.1	4.4	3.2	4.6	5.0	4.5	3.5	6.3	6.9	4.3	6.7	9.3	4.7	5.3	6.4	8.3	9.3	10.4
Beverly-Salem	1.7	2.8	1.2	0.4	1.9	1.1	1.8	1.5	1.6	1.8	3.2	3.9	5.4	2.3	6.3	6.9	8.3	6.1	5.9
Boston Harbor	---	---	---	1.4	1.2	2.0	1.7	2.0	2.1	2.7	2.8	3.0	4.4	4.7	5.0	6.7	6.7	8.0	7.4
Cape Cod Bay	3.9	3.1	3.7	3.1	3.2	2.1	3.9	2.9	3.0	3.3	5.4	6.8	6.8	7.4	10.2	13.6	11.9	13.1	15.5
Outer Cape Cod	11.1	23.0	30.3	26.8	22.3	28.9	16.9	21.4	27.4	24.5	18.3	27.7	26.8	27.3	34.4	34.6	32.9	30.3	34.6
Buzzards Bay	16.0	16.9	32.5	26.6	25.0	25.3	31.0	27.8	29.2	35.0	28.2	28.8	40.9	22.1	26.0	23.6	22.2	32.9	29.5

Table 5. CTHSOD, by state and region, for all ovigerous female American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.024	0.027	0.050	0.038	0.044	0.057	0.049	0.054	0.057	0.073	0.032	0.054	0.092	0.036	0.045	0.047	0.042	0.039	0.039
Cape Ann	0.002	0.011	0.024	0.015	0.016	0.017	0.016	0.010	0.037	0.035	0.024	0.050	0.038	0.024	0.030	0.031	0.056	0.037	0.026
Beverly-Salem	0.011	0.009	0.008	0.003	0.011	0.004	0.010	0.004	0.009	0.005	0.008	0.014	0.020	0.008	0.017	0.015	0.013	0.016	0.012
Boston Harbor	---	---	---	0.009	0.007	0.015	0.012	0.012	0.010	0.028	0.017	0.017	0.026	0.024	0.028	0.032	0.035	0.046	0.032
Cape Cod Bay	0.020	0.025	0.016	0.009	0.015	0.010	0.012	0.009	0.014	0.017	0.028	0.016	0.023	0.022	0.034	0.035	0.027	0.032	0.043
Outer Cape Cod	0.012	0.028	0.040	0.030	0.038	0.032	0.034	0.030	0.043	0.055	0.038	0.076	0.053	0.046	0.081	0.085	0.062	0.045	0.049
Buzzards Bay	0.079	0.053	0.230	0.183	0.193	0.297	0.234	0.289	0.270	0.349	0.073	0.197	0.446	0.110	0.088	0.098	0.083	0.067	0.060

Table 6. CTHAUL, by state and region, for all ovigerous female American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.073	0.078	0.179	0.116	0.133	0.167	0.183	0.189	0.211	0.282	0.140	0.227	0.319	0.143	0.200	0.226	0.219	0.205	0.230
Cape Ann	0.010	0.016	0.038	0.027	0.039	0.047	0.048	0.031	0.096	0.109	0.056	0.088	0.135	0.064	0.085	0.081	0.124	0.097	0.096
Beverly-Salem	0.025	0.033	0.016	0.006	0.033	0.018	0.036	0.021	0.039	0.023	0.049	0.047	0.067	0.048	0.106	0.103	0.136	0.101	0.094
Boston Harbor	---	---	---	0.030	0.025	0.050	0.037	0.038	0.043	0.075	0.064	0.046	0.081	0.088	0.100	0.124	0.132	0.181	0.110
Cape Cod Bay	0.048	0.048	0.040	0.024	0.040	0.031	0.038	0.034	0.039	0.055	0.091	0.056	0.078	0.075	0.128	0.147	0.130	0.110	0.210
Outer Cape Cod	0.081	0.178	0.242	0.170	0.176	0.225	0.157	0.198	0.258	0.342	0.251	0.453	0.317	0.306	0.514	0.561	0.408	0.307	0.330
Buzzards Bay	0.243	0.139	0.828	0.515	0.555	0.748	0.889	0.929	0.953	1.291	0.359	0.847	1.438	0.383	0.343	0.414	0.477	0.527	0.450

Table 7. Estimated fishing pressure index, by state and region, commercial lobster trap catch survey, Massachusetts coastal water, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	86	87	86	86	88	88	89	90	88	87	89	89	90	88	86	86	86	88	88
Cape Ann	91	92	87	89	87	87	88	90	84	81	90	87	83	86	88	85	88	85	80
Beverly-Salem	89	92	94	88	96	96	97	98	96	95	97	98	96	96	95	95	96	94	97
Boston Harbor	---	---	---	93	94	96	96	96	96	95	96	95	96	95	94	95	94	96	97
Cape Cod Bay	90	93	92	94	93	94	92	94	94	93	91	92	94	90	86	88	87	90	87
Outer Cape Cod	46	43	42	38	48	46	54	57	47	50	54	57	60	60	55	54	57	58	46
Buzzards Bay	98	96	96	94	96	97	97	97	95	94	95	97	97	98	96	95	92	96	96

Table 8A. Total instantaneous (Z)* and total annual (A)** mortality estimates (Gulland, 1969) of American lobster by state and region, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	1.58 *	1.72	1.66	1.66	1.76	1.80	1.90	1.86	1.80	1.76	1.90	1.92	1.86	2.02	1.83	1.79	1.87	1.81	1.81
	79% **	82%	81%	81%	83%	84%	85%	84%	83%	83%	85%	85%	84%	87%	84%	83%	85%	84%	84%
Cape Ann	1.65	2.18	1.72	1.92	1.94	2.03	1.85	1.75	1.55	1.39	1.97	1.87	1.51	1.81	1.95	1.90	2.00	1.65	1.48
	81%	89%	82%	85%	86%	87%	84%	83%	79%	75%	86%	85%	78%	84%	86%	85%	86%	81%	77%
Beverly-Salem	1.97	2.15	2.41	2.71	3.64	3.60	3.49	3.31	3.59	2.81	3.49	3.12	2.62	3.34	3.10	2.90	2.46	2.81	3.14
	86%	88%	91%	93%	97%	97%	97%	96%	97%	94%	97%	96%	93%	96%	95%	94%	91%	94%	96%
Boston Harbor	---	---	---	2.52	3.59	2.60	2.77	2.86	2.96	3.00	3.40	3.54	3.26	3.21	2.87	2.65	2.77	2.59	2.86
	---	---	---	92%	97%	93%	94%	94%	95%	95%	97%	97%	96%	96%	94%	93%	94%	92%	94%
Cape Cod Bay	2.53	2.69	2.42	2.52	2.31	2.83	2.26	2.74	2.43	2.46	2.33	2.58	2.60	3.10	2.35	2.09	2.14	2.02	2.14
	92%	93%	91%	92%	90%	94%	90%	94%	91%	91%	90%	92%	93%	95%	90%	88%	88%	87%	88%
Outer Cape Cod	0.43	0.46	0.42	0.33	0.52	0.51	0.80	0.71	0.62	0.63	0.77	0.78	0.87	0.92	0.74	0.73	0.89	0.86	0.58
	35%	37%	34%	28%	41%	40%	55%	51%	46%	47%	54%	54%	58%	60%	52%	52%	59%	58%	44%
Buzzards Bay	3.02	3.00	8.64	3.14	3.55	3.71	3.48	3.18	3.13	2.60	3.50	3.81	3.03	3.58	3.34	2.84	2.74	3.27	3.81
	95%	95%	99%	96%	97%	98%	97%	96%	96%	93%	97%	98%	95%	97%	96%	94%	94%	96%	98%

Table 8B. Total instantaneous (Z)* and total annual (A)** mortality estimates (Beverton and Holt, 1956) of American lobster by state and region, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	1.35	*1.45	1.39	1.41	1.47	1.49	1.54	1.56	1.53	1.50	1.73	1.70	1.79	1.67	1.57	1.54	1.55	1.62	1.62
	74%	**77%	75%	76%	77%	78%	79%	79%	78%	78%	82%	82%	83%	81%	79%	79%	79%	80%	80%
Cape Ann	1.32	1.39	1.35	1.52	1.33	1.32	1.39	1.51	1.27	1.66	1.77	1.57	1.38	1.48	1.62	1.49	1.67	1.52	1.14
	73%	75%	74%	78%	74%	73%	75%	78%	72%	81%	83%	79%	75%	77%	80%	77%	81%	78%	68%
Beverly-Salem	1.59	1.70	1.85	1.78	1.96	1.99	2.16	1.98	2.01	1.83	2.29	2.50	2.23	2.18	2.09	2.11	2.28	2.18	2.08
	80%	82%	84%	83%	86%	86%	88%	86%	87%	84%	90%	92%	89%	89%	88%	88%	90%	89%	88%
Boston Harbor	---	---	---	1.82	1.75	1.92	1.88	1.84	1.94	1.87	2.19	2.14	2.33	2.28	2.09	2.18	2.09	2.63	2.12
	---	---	---	84%	83%	85%	85%	84%	86%	85%	89%	88%	90%	90%	88%	89%	88%	93%	88%
Cape Cod Bay	1.64	1.92	1.72	2.07	1.88	1.92	1.78	1.87	1.97	1.95	1.96	2.01	2.14	1.93	1.65	1.66	1.57	1.65	1.41
	81%	85%	82%	87%	85%	85%	83%	85%	86%	86%	86%	87%	88%	86%	81%	81%	79%	81%	76%
Outer Cape Cod	0.54	0.55	0.53	0.52	0.57	0.55	0.66	0.66	0.62	0.63	0.71	0.72	0.78	0.79	0.72	0.68	0.76	0.76	0.59
	42%	42%	41%	41%	43%	42%	48%	48%	46%	47%	51%	51%	54%	55%	51%	49%	53%	53%	45%
Buzzards Bay	2.97	2.53	2.26	2.21	2.36	2.41	2.36	2.35	2.14	2.27	3.08	2.70	3.11	2.85	2.44	2.37	2.14	2.57	2.42
	95%	92%	90%	89%	91%	91%	91%	94%	88%	90%	95%	93%	96%	94%	91%	91%	88%	92%	91%

Table 9A. Instantaneous fishing mortality estimates (F), by state and region, commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	1.14	1.21	1.17	1.19	1.25	1.28	1.32	1.36	1.36	1.32	1.43	1.44	1.48	1.41	1.32	1.29	1.33	1.36	1.22
Cape Ann	1.33	1.47	1.11	1.33	1.28	1.22	1.30	1.37	1.12	1.04	1.50	1.32	1.14	1.25	1.36	1.28	1.41	1.26	1.07
Beverly-Salem	1.42	1.47	1.64	1.68	1.81	1.93	1.89	2.02	1.95	1.86	2.08	2.16	1.96	1.94	1.88	1.91	2.01	1.87	2.03
Boston Harbor	---	---	---	1.77	1.70	1.80	1.87	1.83	1.94	1.86	2.01	1.97	1.97	1.90	1.85	1.89	1.79	2.13	1.98
Cape Cod Bay	1.53	1.60	1.58	1.73	1.59	1.70	1.56	1.70	1.82	1.72	1.66	1.71	1.85	1.66	1.47	1.44	1.47	1.52	1.40
Outer Cape Cod	0.47	0.48	0.45	0.42	0.47	0.47	0.57	0.53	0.54	0.51	0.59	0.61	0.65	0.68	0.62	0.59	0.65	0.64	0.52
Buzzards Bay	2.32	2.13	1.94	1.80	2.04	2.11	2.08	2.06	1.95	1.97	2.34	2.26	2.39	2.31	2.05	1.98	1.75	2.06	2.14

Table 9B. Instantaneous fishing mortality estimates (F), by state and region, commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.82	0.88	0.86	0.87	0.91	0.95	0.96	0.97	0.96	0.93	0.99	0.99	1.04	0.98	0.93	0.92	0.94	0.97	0.89
Cape Ann	0.82	0.85	0.90	0.92	0.93	0.92	0.92	0.96	0.88	0.76	1.03	0.97	0.86	0.88	0.97	0.89	0.99	0.92	0.75
Beverly-Salem	0.86	1.02	1.05	0.88	1.21	1.23	1.19	1.26	1.25	1.16	1.26	1.32	1.21	1.19	1.19	1.22	1.16	1.18	1.25
Boston Harbor	---	---	---	1.15	1.14	1.10	1.15	1.17	1.22	1.20	1.24	1.21	1.23	1.22	1.17	1.22	1.18	1.34	1.24
Cape Cod Bay	1.04	1.14	1.04	1.19	1.12	1.19	1.09	1.13	1.19	1.17	1.09	1.11	1.19	1.00	0.95	0.91	0.98	1.00	0.99
Outer Cape Cod	0.31	0.34	0.31	0.28	0.34	0.34	0.41	0.38	0.39	0.39	0.44	0.44	0.50	0.51	0.49	0.46	0.52	0.50	0.39
Buzzards Bay	1.10	1.02	1.00	0.94	1.05	1.07	1.04	1.06	1.00	1.00	1.15	1.07	1.14	1.11	0.98	0.99	0.87	1.03	1.07

Table 10. Estimated exploitation rate (u), by state and region, commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.62	0.64	0.63	0.64	0.65	0.66	0.68	0.69	0.69	0.69	0.68	0.69	0.69	0.68	0.66	0.66	0.68	0.67	0.67
Cape Ann	0.74	0.80	0.61	0.68	0.71	0.67	0.70	0.71	0.63	0.51	0.70	0.67	0.62	0.65	0.67	0.66	0.68	0.65	0.64
Beverly-Salem	0.71	0.71	0.75	0.79	0.79	0.83	0.77	0.88	0.76	0.85	0.82	0.79	0.78	0.79	0.79	0.80	0.79	0.76	0.86
Boston Harbor	---	---	---	0.82	0.81	0.80	0.84	0.84	0.86	0.85	0.82	0.81	0.76	0.75	0.78	0.77	0.75	0.75	0.82
Cape Cod Bay	0.75	0.71	0.75	0.73	0.72	0.75	0.73	0.77	0.79	0.76	0.73	0.74	0.76	0.74	0.72	0.70	0.74	0.75	0.75
Outer Cape Cod	0.37	0.37	0.35	0.33	0.36	0.36	0.41	0.38	0.40	0.38	0.42	0.44	0.45	0.47	0.44	0.43	0.45	0.45	0.40
Buzzards Bay	0.74	0.78	0.77	0.72	0.79	0.80	0.80	0.82	0.80	0.78	0.72	0.78	0.74	0.76	0.76	0.76	0.72	0.74	0.80

Table 11. Mean carapace length (mm), by state and region, for all marketable American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	88.5	87.9	88.1	88.7	87.8	87.6	87.5	88.2	88.9	89.0	88.7	88.8	88.4	88.8	89.2	89.2	89.2	88.9	89.6
Cape Ann	88.6	88.3	88.3	87.9	88.4	88.3	88.0	88.3	89.3	90.3	88.4	88.8	89.6	89.6	88.7	89.5	88.7	89.0	90.8
Beverly-Salem	87.6	87.0	86.6	86.9	86.2	86.2	85.8	87.1	87.7	88.3	87.5	87.2	87.5	87.8	88.0	87.9	87.5	87.7	87.6
Boston Harbor	---	---	---	86.8	86.9	86.4	86.6	87.5	88.0	88.1	87.8	87.9	87.5	87.5	88.0	87.7	87.8	87.0	87.5
Cape Cod Bay	87.2	86.4	86.9	86.1	86.4	86.3	86.7	87.3	87.7	87.7	88.1	88.2	87.7	88.3	89.2	89.0	89.4	89.0	89.4
Outer Cape Cod	98.2	97.5	97.4	99.7	97.0	96.3	94.6	95.2	96.5	96.1	95.3	95.2	93.8	94.2	94.2	94.9	93.9	94.1	96.6
Buzzards Bay	84.7	85.2	85.7	85.8	85.2	85.3	85.3	86.1	87.4	87.0	86.4	86.9	86.5	86.5	87.4	87.3	87.8	87.0	87.0

Table 12. Mean carapace length (mm), by state and region for all sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	75.8	76.3	76.2	76.1	76.3	76.1	76.1	76.3	77.5	77.6	76.7	76.2	76.9	77.5	77.8	78.2	77.8	77.2	77.4
Cape Ann	78.0	77.7	77.5	77.3	77.6	77.1	75.9	77.0	78.3	78.8	78.7	77.9	78.3	78.0	77.2	77.3	77.3	77.8	76.8
Beverly-Salem	74.3	76.5	74.9	76.1	75.9	74.7	74.7	74.5	76.4	76.1	73.4	73.5	75.1	75.8	76.2	76.8	76.0	74.9	75.4
Boston Harbor	---	---	---	77.1	76.9	76.9	76.5	75.6	76.8	77.4	75.4	74.6	75.3	76.0	77.3	76.9	76.3	72.7	74.9
Cape Cod Bay	76.6	76.4	76.7	75.6	76.1	76.2	75.6	76.9	77.9	77.8	77.4	76.8	76.7	78.6	78.6	79.8	79.1	79.3	78.9
Outer Cape Cod	75.9	76.2	77.1	75.1	76.6	75.9	77.0	77.1	76.8	78.8	78.8	79.0	79.4	78.3	80.0	79.9	80.0	79.6	79.6
Buzzards Bay	75.8	75.5	76.8	76.4	76.1	76.0	76.6	76.3	77.7	77.4	76.6	77.1	78.3	77.6	77.7	77.5	78.2	78.9	78.7

Table 13. Mean carapace length (mm) of all ovigerous female American lobster, by state and region, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	88.5	87.6	88.6	87.4	87.9	88.1	87.1	87.2	88.5	88.0	86.0	85.5	85.3	86.3	86.6	86.7	86.7	86.4	88.1
Cape Ann	109.0	100.3	94.3	90.5	93.8	95.0	91.6	94.0	100.4	95.1	91.7	91.9	91.0	92.7	93.3	91.5	90.7	91.3	94.7
Beverly-Salem	80.5	84.5	85.8	83.5	85.9	83.5	81.8	83.0	85.2	85.5	83.8	81.6	82.6	83.1	83.3	83.0	82.5	83.0	83.0
Boston Harbor	---	---	---	82.1	84.0	81.3	82.3	83.7	83.0	83.8	82.0	82.0	80.8	80.9	81.5	82.4	82.2	79.8	81.2
Cape Cod Bay	86.4	83.8	85.5	84.4	85.2	86.8	87.0	84.7	86.1	85.0	83.9	84.1	83.0	84.8	85.2	85.7	85.8	86.1	87.3
Outer Cape Cod	109.8	106.1	108.0	107.1	106.9	107.3	102.5	105.2	105.4	104.6	101.9	99.2	100.7	100.0	100.6	99.6	98.9	98.3	104.7
Buzzards Bay	78.1	79.6	81.6	83.0	80.1	79.4	80.2	80.6	81.3	80.8	79.8	79.9	81.0	81.5	81.8	82.6	84.0	82.7	82.8

Table 14. Cull rate (percent), by state and region, for all American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal water, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	10.0	10.8	10.7	14.8	18.1	20.9	17.0	18.2	19.2	18.6	18.1	18.5	20.3	21.9	22.0	17.1	20.4	20.9	20.8
Cape Ann	10.0	9.8	10.5	11.5	23.9	25.3	20.2	21.2	16.7	16.7	19.7	18.2	19.2	17.1	19.6	18.0	18.9	17.6	16.7
Beverly-Salem	8.3	8.6	10.2	20.9	23.0	30.0	24.1	26.3	28.6	27.3	28.9	22.7	28.3	30.8	25.1	24.6	23.4	25.1	23.6
Boston Harbor	---	---	---	13.3	19.3	19.1	16.9	16.3	13.8	14.7	13.5	17.2	23.4	23.0	22.2	17.8	19.8	17.4	17.7
Cape Cod Bay	11.1	10.7	10.9	15.6	18.3	21.6	16.2	17.4	22.8	20.5	18.9	18.3	18.1	19.4	21.5	16.2	23.6	25.6	24.6
Outer Cape Cod	5.7	11.3	8.9	13.0	13.4	16.1	12.6	15.0	14.0	15.5	13.2	15.7	17.3	20.1	19.0	13.3	14.5	14.2	14.9
Buzzards Bay	13.5	14.7	12.4	12.4	13.4	14.6	15.1	15.6	12.6	13.6	13.9	19.3	20.5	24.0	24.4	15.0	15.6	16.2	18.5

Table 15. Cull rate (percent), by state and region, for all legal-sized American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	8.1	9.7	9.2	12.7	14.8	17.0	14.7	15.7	14.9	15.4	15.6	17.1	17.4	21.2	20.9	15.1	19.3	19.5	19.1
Cape Ann	10.7	9.6	7.5	10.4	19.4	20.3	18.0	19.3	13.9	13.7	16.8	18.3	16.3	16.5	16.2	14.7	15.7	16.3	13.7
Beverly-Salem	4.3	7.7	7.4	15.5	19.3	22.1	17.1	21.4	18.7	25.6	22.8	19.9	24.6	25.4	18.8	20.2	17.2	20.0	18.2
Boston Harbor	---	---	---	10.1	16.2	15.8	12.9	13.1	9.9	9.9	12.3	14.0	17.5	18.0	20.1	17.3	17.3	14.2	14.0
Cape Cod Bay	9.3	9.3	10.0	13.2	14.5	18.1	15.0	15.6	12.0	16.3	17.8	16.8	16.3	21.7	23.3	16.0	25.3	26.2	24.3
Outer Cape Cod	5.3	10.3	8.1	13.3	12.5	14.9	13.1	14.3	13.3	14.1	12.8	15.3	16.4	19.9	18.3	12.6	13.6	13.5	14.5
Buzzards Bay	16.1	13.2	12.7	12.3	13.8	13.6	15.2	14.1	12.6	12.6	11.5	22.2	18.9	23.5	22.6	12.9	15.3	14.4	17.8

Table 16. Cull rate (percent), by state and region, for marketable American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	8.2	9.9	9.2	13.2	16.2	17.6	14.7	16.0	15.2	15.6	16.1	17.6	17.6	21.9	21.5	15.6	19.2	19.7	19.6
Cape Ann	10.8	9.8	7.3	10.5	20.9	20.7	18.4	19.9	14.0	14.2	16.8	18.1	13.7	16.7	16.0	14.5	15.3	16.3	14.1
Beverly-Salem	4.4	8.0	7.4	15.6	18.5	22.2	17.2	21.3	18.9	23.8	23.1	20.0	24.7	25.5	19.0	20.0	17.0	20.1	18.0
Boston Harbor	---	---	---	10.2	16.2	15.7	12.8	13.1	9.9	9.9	12.4	14.0	17.5	18.0	20.1	17.4	17.4	14.5	14.2
Cape Cod Bay	9.3	9.3	10.0	13.2	15.9	18.2	14.8	15.6	19.1	16.2	17.8	16.7	16.2	22.3	23.3	16.6	25.5	26.1	24.7
Outer Cape Cod	5.3	10.9	8.6	14.8	12.9	16.8	13.2	14.9	13.9	14.6	14.1	16.8	17.3	21.6	20.4	14.0	13.7	14.0	15.6
Buzzards Bay	16.9	13.1	12.3	12.6	15.4	14.1	15.4	14.7	13.0	12.4	11.7	22.5	19.4	23.7	23.0	13.1	14.5	15.1	17.7

Table 17. Cull rate (percent), by state and region, for sub-legal American lobster, sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	11.2	11.5	11.6	16.1	20.2	23.2	18.2	19.6	21.1	20.2	19.2	19.3	21.8	22.5	22.9	18.5	21.3	21.9	22.7
Cape Ann	8.0	10.6	12.6	12.2	26.9	28.7	21.5	22.1	17.9	18.3	21.0	18.2	20.9	17.4	21.2	20.6	20.8	18.4	20.5
Beverly-Salem	10.0	9.0	11.2	22.3	24.0	31.8	25.3	28.6	30.8	29.2	31.6	23.5	29.5	33.8	28.2	26.7	25.5	27.2	27.5
Boston Harbor	---	---	---	14.5	20.5	20.0	18.0	18.0	15.2	16.4	13.9	18.3	25.3	24.7	22.8	17.9	20.6	18.0	19.6
Cape Cod Bay	11.9	11.3	11.4	17.0	20.2	23.4	16.8	18.3	24.0	21.8	19.2	19.0	18.8	17.8	20.1	16.3	22.2	25.2	25.1
Outer Cape Cod	7.8	17.9	13.5	11.7	18.6	22.8	11.0	16.9	17.1	20.7	14.3	17.1	20.2	21.3	21.8	15.7	17.9	16.6	18.3
Buzzards Bay	12.7	15.2	12.2	12.4	13.3	14.9	15.0	16.2	12.6	13.9	14.5	18.3	21.0	24.2	25.6	16.4	15.8	17.4	19.1

Table 18. Percent trap mortality by state and region for all American lobster sampled during commercial lobster trap catch survey, Massachusetts coastal waters, 1981-1999.

	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999
State	0.15	0.04	0.22	0.15	0.18	0.20	0.10	0.15	0.12	0.17	0.37	0.08	0.13	0.22	0.21	0.18	0.10	0.15	0.18
Cape Ann	0.00	0.00	0.09	0.27	0.03	0.16	0.00	0.03	0.13	0.09	0.48	0.10	0.11	0.14	0.28	0.17	0.04	0.13	0.40
Beverly-Salem	0.00	0.00	0.00	0.00	0.04	0.22	0.03	0.19	0.14	0.29	0.41	0.13	0.19	0.13	0.74	0.49	0.37	0.31	0.39
Boston Harbor	---	---	---	0.00	0.03	0.03	0.23	0.09	0.03	0.04	0.01	0.03	0.06	0.04	0.01	0.04	0.08	0.08	0.19
Cape Cod Bay	0.00	0.02	0.03	0.00	0.00	0.02	0.15	0.00	0.02	0.05	0.02	0.02	0.02	0.00	0.03	0.03	0.03	0.08	0.04
Outer Cape Cod	0.46	0.22	0.23	0.48	0.40	0.85	0.27	0.66	0.47	0.62	0.35	0.24	0.30	0.58	0.38	0.43	0.21	0.25	0.25
Buzzards Bay	0.62	0.00	1.13	0.43	0.76	0.25	0.01	0.18	0.11	0.18	1.74	0.10	0.29	0.71	0.16	0.21	0.00	0.19	0.20